

NUMERICAL SOLUTION OF A PARABOLIC PROBLEM ARISING IN FINANCE

MARIE-NOELLE LE ROUX

ABSTRACT. In this paper, we study a parabolic system of three equations which permits to solve an optimal replication problem in incomplete markets. We obtain existence and uniqueness of the solution in suitable Sobolev spaces and propose a numerical method to compute the optimal strategy.

1. INTRODUCTION

We study here a parabolic system arising in the resolution of an optimal replication problem in incomplete markets. Given a European derivative security with an arbitrary payoff function, the optimal replication problem is to find a dynamic portfolio strategy, that is self-financing and comes as close as possible to the payoff at maturity date T . In complete markets, such a dynamic-hedging strategy exists: the payoff of a European option can be replicated exactly; it is the Black-Scholes model (1973) [2]. In [1], Bertsimas, Kogan and Lo propose a solution approach for this problem in incomplete markets..

At time $\tau = 0$, consider a portfolio of stocks and riskless bonds at a cost V_0 and denote by $\theta(\tau)$, $B(\tau)$, $V(\tau)$ the number of shares of the stock held, the value of bonds held and the market value of the portfolio at time τ . Hence, $V(\tau) = \theta(\tau)P(\tau) + B(\tau)$, $0 \leq \tau \leq T$.

If we note σ the volatility and F the payoff function, the value function J is defined by:

$$J(\tau, V, P, \sigma) = \min_{\theta(s), s \geq \tau} E(((V(T) - F(P(T), \sigma(T)))^2 / (V(\tau), P(\tau), \sigma(\tau))).$$

The replication error $\epsilon(V_0)$ is $(J(0, V, P, \sigma))^{1/2}$ and it can be minimized with respect to the initial wealth V_0 to yield the least-cost optimal-replication strategy and the minimum replication error ϵ^* is $\epsilon^* = \min_{V_0} \epsilon(V_0)$.

In [1], it has been proved that the value function J is quadratic in V : $J = a(V - b)^2 + c$ and the coefficients a , b , c satisfy the following system of partial differential equations:

$$(1.1) \quad \frac{\partial a}{\partial \tau} = -\frac{k^2 \sigma^2}{2} \frac{\partial^2 a}{\partial \sigma^2} - g_1(\sigma) \frac{\partial a}{\partial \sigma} + \rho^2 k^2 \frac{\sigma^2}{a} \left(\frac{\partial a}{\partial \sigma} \right)^2 + a f^2(\sigma),$$

$$\frac{\partial b}{\partial \tau} = -\frac{k^2 \sigma^2}{2} \frac{\partial^2 b}{\partial \sigma^2} - \frac{\sigma^2 P^2}{2} \frac{\partial^2 b}{\partial P^2} - \rho k \sigma^2 P \frac{\partial^2 b}{\partial \sigma \partial P} - g_2(\sigma) \frac{\partial b}{\partial \sigma}$$

Key words and phrases. mathematical model, nonlinear parabolic problem, finite elements method.

$$(1.2) \quad - (1 - \rho^2) k^2 \frac{\sigma^2}{a} \frac{\partial a}{\partial \sigma} \frac{\partial b}{\partial \sigma},$$

$$\frac{\partial c}{\partial \tau} = -\frac{k^2 \sigma^2}{2} \frac{\partial^2 c}{\partial \sigma^2} - \frac{\sigma^2 P^2}{2} \frac{\partial^2 c}{\partial P^2} - \rho k \sigma^2 P \frac{\partial^2 c}{\partial \sigma \partial P} - g(\sigma) \frac{\partial c}{\partial \sigma} - \sigma f(\sigma) P \frac{\partial c}{\partial P}$$

$$(1.3) \quad - (1 - \rho^2) k^2 \sigma^2 a \left(\frac{\partial b}{\partial \sigma} \right)^2,$$

where $k > 0$, $\rho \in [-1, +1]$, (ρ is a correlation coefficient)

$g(\sigma) = -\delta \sigma (\sigma - \sigma_1)$ ($\delta > 0$ and $\sigma_1 \in]0, 1[$)

$g_1(\sigma) = g(\sigma) - 2\rho k \sigma f(\sigma)$, $g_2(\sigma) = g(\sigma) - \rho k \sigma f(\sigma)$

and $f(\sigma) = \begin{cases} \frac{\mu}{\sigma_0} & \text{if } \sigma \leq \sigma_0 \\ \frac{\mu}{\sigma} & \text{if } \sigma \geq \sigma_0 \end{cases}$, $\mu > 0$, (μ is the drift).

Remark 1.1. *The function f has been modified near 0 in order to be bounded and assure the existence of a solution.*

The conditions at the time expiry T are given by: $a(T) = 1$, $b(T) = F(P, \sigma)$, $c(T) = 0$.

Under the optimal replication strategy θ^* , the minimum replication error as a function of the initial wealth V_0 is $(J(0))^{\frac{1}{2}} = (a(0)(V_0 - b(0))^2 + c(0))^{\frac{1}{2}}$, hence the initial wealth that minimizes the replication error is $V_0^* = b(0)$ the minimal replication error over all V_0 is $\epsilon^* = \sqrt{c(0)}$ and the least-cost optimal strategy at $\tau = 0$ is $\theta^*(0) = \frac{\partial b}{\partial P}(0) + \frac{\rho k}{P} \frac{\partial b}{\partial \sigma}(0)$.

Remark 1.2. *Exact replication is possible when $k^2(1 - \rho^2) = 0$ and this corresponds to the following cases:*

- *Volatility is a deterministic function of time.*
- *The Brownian motions driving stocks prices and volatility are perfectly correlated.*

In this paper, we propose a numerical method to compute the solution of equations (1.1), (1.2), (1.3) and then obtain the minimal replication error and the least-cost optimal replication strategy.

To obtain a forward problem, we change the sense of time; we note $t = T - \tau$. In order to avoid the function a at the denominator, we make the change of unknown $u_1 = \ln(a)$. We also replace σ by x , P by y , b by u_2 and c by u_3 .

The preceding system becomes:

$$(1.4) \quad \frac{\partial u_1}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 u_1}{\partial x^2} - g_1(x) \frac{\partial u_1}{\partial x} + k^2 \left(\rho^2 - \frac{1}{2} \right) x^2 \left(\frac{\partial u_1}{\partial x} \right)^2 + f^2(x) = 0,$$

$$(1.5) \quad \begin{aligned} & \frac{\partial u_2}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 u_2}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 u_2}{\partial y^2} - \rho k x^2 y \frac{\partial^2 u_2}{\partial x \partial y} - g_2(x) \frac{\partial u_2}{\partial x} \\ & - (1 - \rho^2) k^2 x^2 \frac{\partial u_1}{\partial x} \frac{\partial u_2}{\partial x} = 0, \end{aligned}$$

$$(1.6) \quad \begin{aligned} & \frac{\partial u_3}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 u_3}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 u_3}{\partial y^2} - \rho k x^2 y \frac{\partial^2 u_3}{\partial x \partial y} - g(x) \frac{\partial u_3}{\partial x} - x f(x) y \frac{\partial u_3}{\partial y} \\ & - (1 - \rho^2) k^2 x^2 \exp(u_1) \left(\frac{\partial u_2}{\partial x} \right)^2 = 0, \end{aligned}$$

with the initial conditions:

$u_1(0) = 0$; $u_2(0) = F(x, y)$; $u_3(0) = 0$. (F is the payoff function).

The outline of the paper is as follows:

In section 2, we solve (1.4). The different derivative terms will be treated separately in order to obtain the L^∞ - stability of the scheme. We prove the convergence of the numerical solution towards a weak solution of the problem. Besides the uniqueness of this weak solution is obtained.

In sections 3 and 4, we study (1.5), (1.6). We use a change of unknown which lead to a variationnel formulation and obtain the existence of a unique solution in suitable weighted Sobolev spaces. These equations are discretized by using a backward Euler method in time and a finite element method in space. Numerical results are presented.

2. COMPUTATION OF u_1

2.1. Definition of the numerical solution. In order to solve (1.4), we use suitable weighted Sobolev spaces, such that no boundary condition is needed in 0 and the function u_1 has the correct behaviour at infinity.

To simplify notation, we denote:

$$F_1(x) = f^2(x), \quad x > 0, \quad \lambda = k^2 \left(\rho^2 - \frac{1}{2} \right),$$

this coefficient may be positive or negative since ρ is a correlation factor and then ρ lies in $[-1, +1]$. The equation (1.4) becomes:

$$(2.1) \quad \frac{\partial u_1}{\partial t} - \frac{1}{2} k^2 x^2 \frac{\partial^2 u_1}{\partial x^2} - g_1(x) \frac{\partial u_1}{\partial x} + \lambda x^2 \left(\frac{\partial u_1}{\partial x} \right)^2 + F_1(x) = 0.$$

We will make the following assumptions on the functions f and g_1 :

- $f \in W^{1,\infty}(\mathbb{R}^+)$, $xf' \in L^\infty(\mathbb{R}^+)$.

- F_1 is a nonincreasing function, F_1' is a bounded variation function; if we denote $\hat{F}$ the function defined by $\hat{F}(x) = xF_1'(x)$, $x > 0$, we get $\hat{F} \in L^\infty(\mathbb{R}^+)$.

- g_1 may be written $g_1(x) = x\phi(x)$ with $\phi(x) = -\delta(x - \sigma_1) - 2\rho kf(x)$, $\delta > 0$; so, there exists $\sigma_2 > 0$ such that ϕ is negative and nonincreasing on $[\sigma_2, +\infty[$ and bounded on $[0, \sigma_2]$.

We define the two constants c_1 and c_2 by

$$(2.2) \quad c_1 = \sup_{x \in [0, \sigma_2]} |g_1'(x)|, \quad c_2 = \sup_{x \in [0, \sigma_2]} |\phi(x)|$$

We denote by Δt_n the time increment between the levels t_n and t_{n+1} , $n \geq 0$ and by u_{1h}^n the approximate solution at the time level t_n . This solution will be in a finite-dimensional space V_{1h} which will be defined below.

The solution u_{1h}^{n+1} at the time level t_{n+1} is computed in two steps: knowing u_{1h}^n , we compute $u_{1h}^{n+\frac{1}{2}}$, approximate solution of

$$(2.3) \quad \frac{\partial u_1}{\partial t} + \lambda x^2 \left(\frac{\partial u_1}{\partial x} \right)^2 = 0$$

obtained by using an explicit upwind scheme. Then starting with this intermediate value, we use a backward Euler method in time to compute u_{1h}^{n+1} ; the second order term in (2.1) is discretized by using a P_1 -finite element method [4] and the linear first order term by an implicit upwind scheme [6] in order to get the L^∞ -stability of the scheme.

In order to define the finite-dimensional space V_{1h} , we first study the parabolic problem:

$$(2.4) \quad \frac{\partial u_1}{\partial t} - \frac{1}{2}k^2x^2 \frac{\partial^2 u_1}{\partial x^2} + F_1(x) = 0$$

to obtain a variational formulation in weighted Sobolev spaces.

2.1.1. Variational formulation of (2.4).

Let us consider the two spaces:

$$H_1 = \left\{ v \in \mathcal{D}'(\mathbb{R}^+) / \frac{v}{1+x} \in L^2(\mathbb{R}^+) \right\}$$

and

$$V_1 = \left\{ v \in \mathcal{D}'(\mathbb{R}^+) / \frac{v}{1+x} \in L^2(\mathbb{R}^+), \frac{xv'}{1+x} \in L^2(\mathbb{R}^+) \right\}.$$

The space H_1 is equipped with the following scalar product:

$$\forall v, w \in H_1, (v, w)_1 = \int_0^{+\infty} \frac{v(x)w(x)}{(1+x)^2} dx \text{ and the associated norm.}$$

The space V_1 with the norm

$$\|v\|_{V_1} = \left(\int_0^\infty \left(\frac{v^2(x)}{(1+x)^2} + \frac{x^2}{(1+x)^2} \left(\frac{dv}{dx} \right)^2 \right) dx \right)^{\frac{1}{2}}$$

is a Hilbert space and $\mathcal{D}(\mathbb{R}^+)$ is dense in V_1 [3].

We define on $V_1 \times V_1$ the bilinear form:

$$\forall v, w \in V_1, a(v, w) = \frac{1}{2} k^2 \int_0^{+\infty} \frac{dv}{dx} \frac{dw}{dx} \left(w \frac{x^2}{(1+x)^2} \right) dx$$

or

$$a(v, w) = \frac{1}{2} k^2 \int_0^{+\infty} \frac{x^2}{(1+x)^2} \frac{dv}{dx} \frac{dw}{dx} dx + k^2 \int_0^{+\infty} \frac{x}{(1+x)^3} \frac{dv}{dx} w dx.$$

This bilinear form is continue on $V_1 \times V_1$ and we have the equality

$$a(v, v) = \frac{k^2}{2} \int_0^{+\infty} \frac{x^2}{(1+x)^2} \left(\frac{dv}{dx} \right)^2 dx - \frac{k^2}{2} \int_0^{+\infty} \frac{1-2x}{(1+x)^4} v^2 dx,$$

then we get:

$$\forall v \in V_1, a(v, v) \geq \frac{k^2}{2} \|v\|_{V_1}^2 - k^2 \|v\|_{H_1}^2.$$

Since the function F_1 is in H_1 , the following variational problem:

Find $u_1 \in L^2(0, T; V_1) \cap C(0, T; H_1)$ such that:

$$(2.5) \quad \begin{cases} \left(\frac{\partial u_1}{\partial t}, v \right)_1 + a(u_1, v) = -(F_1, v)_1, \quad \forall v \in V_1. \\ u_1(0) = 0 \end{cases}$$

has a unique solution [5].

2.1.2. Approximation of (2.5).

The finite-dimensional space V_{1h} will be a subspace of V_1 defined in the following way:

Let $(x_i)_{0 \leq i \leq N}$ an increasing sequence ($x_0 = 0$). We denote $h_i = x_i - x_{i-1}$, $I_i = (x_{i-1}, x_i)$, $1 \leq i \leq N$, $I_{N+1} = (x_N, +\infty)$

$$V_{1h} = \{ v_h \in C^0(\mathbb{R}^+) / v_h|_{I_i} \in P_1, 1 \leq i \leq N, v_h|_{I_{N+1}} \in P_0 \}.$$

The variable x is the volatility which lies, in practice, in $]0, 1[$, so, we may use a constant space step h on $(0, 1)$ and an increasing sequence (h_i) for $x \geq 1$ in order that the number of nodes is not too important.

If $v_h \in V_{1h}$, we denote $v_i = v_h(x_i)$.

We define on V_{1h} an approximate scalar product:

$$(v_h, w_h)_h = \frac{h_1}{2} v_0 w_0 + \sum_{i=1}^{N-1} \frac{h_i + h_{i+1}}{2} \frac{1}{(1+x_i)^2} v_i w_i + v_N w_N \left(\frac{h_N}{2(1+x_N)^2} + \frac{1}{1+x_N} \right)$$

obtained by using the trapezoid method on each interval I_i , $1 \leq i \leq N$; the last integral being computed exactly.

We also define the Lagrange interpolate $\pi_h F_1$ of F_1 by:

$\pi_h F_1 \in V_{1h}$ and $\pi_h F_1(x_i) = F_1(x_i) = F_{1i}$, $1 \leq i \leq N$.

The approximate solution of (2.5) at the time level t_{n+1} is the solution of :

$$(u_{1h}^{n+1}, v_h)_h + \Delta t_n a(u_{1h}^{n+1}, v_h) = (u_{1h}^n, v_h)_h - \Delta t_n (\pi_h F_1, v_h)_h, \quad \forall v_h \in V_{1h},$$

$$u_{1h}^0 = 0.$$

This may be written:

$$u_{10}^{n+1} = u_{10}^n - \Delta t_n F_{10},$$

$$u_{1i}^{n+1} + \Delta t_n \alpha_i \left(\left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) u_{1i}^{n+1} - \frac{1}{h_i} u_{1,i-1}^{n+1} - \frac{1}{h_{i+1}} u_{1,i+1}^{n+1} \right) = u_{1i}^n - \Delta t_n F_{1i},$$

$$u_{1N}^{n+1} + \frac{\Delta t_n}{h_N} \alpha_N (u_{1N}^{n+1} - u_{1,N-1}^{n+1}) = u_{1N}^n - \Delta t_n F_{1N},$$

$$\text{with } \alpha_i = \frac{k^2 x_i^2}{h_i + h_{i+1}}, \quad 1 \leq i \leq N-1, \quad \alpha_N = \frac{k^2 x_N^2}{h_N + 2(1 + x_N)}.$$

2.1.3. Approximation of the first order terms. We compute now an approximate solution of (2.3) by using an explicit upwind scheme.

Let us denote by v_h^n the derivative of u_{1h}^n

$$v_h^n | I_i = v_i^n = \frac{u_{1i}^n - u_{1,i-1}^n}{h_i}, \quad 1 \leq i \leq N, \quad v_h^n | I_{N+1} = v_{N+1}^n = 0.$$

We set $v_0^n = 0$.

We shall prove below that the function v_h^n is positive and the function u_{1h}^n is negative; we define $u_{1h}^{n+\frac{1}{2}} \in V_{1h}$ by:

$$(2.6) \quad u_{1i}^{n+\frac{1}{2}} = u_{1i}^n - \lambda \Delta t_n x_i^2 (v_i^n)^2 \quad \text{if } \lambda > 0,$$

$$(2.7) \quad u_{1i}^{n+\frac{1}{2}} = u_{1i}^n - \lambda \Delta t_n x_{i+1}^2 (v_{i+1}^n)^2 \quad \text{if } \lambda < 0,$$

$0 \leq i \leq N$.

For the linear first order term, since the function g_1 is not bounded, we use an implicit scheme, which will be decentered in order to get a monotone matrix.

Finally, the solution $u_{1h}^{n+1} \in V_{1h}$ of (2.1) is defined by:

$$\begin{aligned}
 u_{10}^{n+1} &= u_{10}^{n+\frac{1}{2}} - \Delta t_n F_{10}, \\
 u_{1i}^{n+1} + \Delta t_n \left(\alpha_i \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) + \gamma_i \left(\frac{1 - \delta_i}{h_{i+1}} - \frac{\delta_i}{h_i} \right) \right) u_{1i}^{n+1} \\
 &\quad - \frac{\Delta t_n}{h_i} (\alpha_i - \gamma_i \delta_i) u_{1,i-1}^{n+1} - \frac{\Delta t_n}{h_{i+1}} (\alpha_i + \gamma_i (1 - \delta_i)) u_{1,i+1}^{n+1} \\
 &= u_{1i}^{n+\frac{1}{2}} - \Delta t_n F_{1i},
 \end{aligned}
 \tag{2.8}$$

$$\begin{aligned}
 u_{1N}^{n+1} + \frac{\Delta t_n}{h_N} (\alpha_N - \gamma_N) (u_{1N}^{n+1} - u_{1,N-1}^{n+1}) \\
 = u_{1N}^{n+\frac{1}{2}} - \Delta t_n F_{1N},
 \end{aligned}$$

with $\gamma_i = g_1(x_i)$, $1 \leq i \leq N$,

$$\delta_i = \begin{cases} 0 & \text{if } \gamma_i \geq 0, \\ 1 & \text{if } \gamma_i < 0. \end{cases}$$

Since g_1 is negative for $x \geq \sigma_2$, we get $\delta_i = 1$ if i is large enough.

The preceding equations may be written by using the derivative v_h^n :

$$u_{1i}^{n+1} + \Delta t_n ((\alpha_i - \delta_i \gamma_i) v_i^{n+1} - (\alpha_i + (1 - \delta_i) \gamma_i) v_{i+1}^{n+1}) = u_{1i}^{n+\frac{1}{2}} - \Delta t_n F_{1i},$$

$$0 \leq i \leq N$$

2.2. Properties of the scheme. We prove that under a stability condition, the approximate solution u_{1h}^n is negative and its derivative v_h^n is positive.

Let us denote U_{1h}^n the vector of $\mathbb{R}^{N+1}$ of components (u_{1i}^n) , $0 \leq i \leq N$ and F_{1h} the vector of components (F_{1i}) , $0 \leq i \leq N$.

The numerical scheme (2.8) may be written:

$$(I + \Delta t_n A_h) U_{1h}^{n+1} = U_{1h}^{n+\frac{1}{2}} - \Delta t_n F_{1h}
 \tag{2.9}$$

where the matrix A_h is tridiagonal and monotone.

From (2.6), (2.7) and (2.8), we get immediately the equations satisfied by v_h^n :

If $v_h^{n+\frac{1}{2}}$ denotes the derivative of $u_h^{n+\frac{1}{2}}$, we obtain for $1 \leq i \leq N$

$$(2.10) \quad v_i^{n+\frac{1}{2}} = v_i^n - \lambda \frac{\Delta t_n}{h_i} (x_i^2 (v_i^n)^2 - x_{i-1}^2 (v_{i-1}^n)^2) \quad \text{if } \lambda > 0,$$

$$(2.11) \quad v_i^{n+\frac{1}{2}} = v_i^n - \lambda \frac{\Delta t_n}{h_i} (x_{i+1}^2 (v_{i+1}^n)^2 - x_i^2 (v_i^n)^2) \quad \text{if } \lambda < 0,$$

and v_h^{n+1} satisfies:

$$(2.12) \quad \begin{aligned} & v_i^{n+1} + \frac{\Delta t_n}{h_i} (\alpha_i + \alpha_{i-1} - \delta_i \gamma_i + (1 - \delta_{i-1}) \gamma_{i-1}) v_i^{n+1} \\ & - \frac{\Delta t_n}{h_i} (\alpha_{i-1} - \delta_{i-1} \gamma_{i-1}) v_{i-1}^{n+1} - \frac{\Delta t_n}{h_i} (\alpha_i + (1 - \delta_i) \gamma_i) v_{i+1}^{n+1} \\ & = v_i^{n+\frac{1}{2}} - \frac{\Delta t_n}{h_i} (F_{1i} - F_{1,i-1}) \end{aligned}$$

for $1 \leq i \leq N$
which may be written:

$$(2.13) \quad (I + \Delta t_n B_h) V_h^{n+1} = V_h^{n+\frac{1}{2}} - \Delta t_n G_{1h},$$

where V_h^n is the vector of $\mathbb{R}^N$ of components (v_i^n) , $1 \leq i \leq N$, B_h is a tridiagonal matrix $(N \times N)$, G_{1h} is the vector of $\mathbb{R}^N$ of components $\frac{F_{1i} - F_{1,i-1}}{h_i}$, $1 \leq i \leq N$.

Proposition 2.1. *If the following stability condition*

$$(2.14) \quad \sup_{i \geq 1} \left| \lambda \frac{\Delta t_n}{h_i} x_i^2 v_i^n \right| \leq 1$$

is satisfied and

$$(2.15) \quad \Delta t_n c_1 < 1,$$

then the function v_h^n is nonnegative for $n \geq 0$.

Proof: We can rewrite (2.10) as:

$$(2.16) \quad v_i^{n+\frac{1}{2}} = v_i^n \left(1 - \lambda \frac{\Delta t_n}{h_i} x_i^2 v_i^n \right) + \lambda \frac{\Delta t_n}{h_i} x_{i-1}^2 (v_{i-1}^n)^2 \quad \text{si } \lambda > 0$$

and we have an analogous formula for $\lambda < 0$.

If (2.14) is satisfied, we get immediately that $v_h^n \geq 0$ implies $v_h^{n+\frac{1}{2}} \geq 0$. Since F_1 is decreasing, the vector $V_h^{n+\frac{1}{2}} - \Delta t_n G_{1h}$ is nonnegative and the vector V_h^{n+1} will be nonnegative if $I + \Delta t_n B_h$ is a monotone matrix; this will be true if $1 - \frac{\Delta t_n}{h_i}(\gamma_i - \gamma_{i-1}) > 0$, for $1 \leq i \leq N$. Since g_1 is a decreasing function for $x \geq \sigma_2$, this condition is satisfied for i large enough and if $x \leq \sigma_2$, we get $1 - \frac{\Delta t_n}{h_i}(\gamma_i - \gamma_{i-1}) \geq 1 - c_1 \Delta t_n > 0$ from (2.15).

We deduce immediately the following results:

Proposition 2.2. *Under the hypotheses of proposition 2.1, the numerical solution u_{1h}^n satisfies: $u_{1h}^n \leq 0$ for $n \geq 0$.*

Proof: We can rewrite (2.6) as

$$(2.17) \quad u_{1i}^{n+\frac{1}{2}} = u_{1i}^n \left(1 - \lambda \frac{\Delta t_n}{h_i} x_i^2 v_i^n \right) + \lambda \frac{\Delta t_n}{h_i} x_i^2 v_i^n u_{1,i-1}^n \text{ if } \lambda > 0$$

and we have an analogous equality for $\lambda < 0$. From proposition (2.1), the function v_h^n is nonnegative, hence if $u_{1h}^n \leq 0$, we get $u_{1h}^{n+\frac{1}{2}} \leq 0$.

Since $I + \Delta t_n A_h$ is a monotone matrix and $F_1 \geq 0$, we deduce that $u_{1h}^{n+1} \leq 0$.

Proposition 2.3. *Under the hypotheses of proposition 2.1, the numerical solution satisfies*

$$\|u_{1h}^n\|_{L^\infty(\mathbb{R}^+)} \leq t_n F_1(0)$$

for $n \geq 0$.

Proof: We get immediately from (2.17): $\left\| u_{1h}^{n+\frac{1}{2}} \right\|_{L^\infty(\mathbb{R}^+)} \leq \|u_{1h}^n\|_{L^\infty(\mathbb{R}^+)}$ and since A_h is a monotone matrix, it follows from (2.9) that

$$\|u_{1h}^{n+1}\|_{L^\infty(\mathbb{R}^+)} \leq \left\| u_{1h}^{n+\frac{1}{2}} \right\|_{L^\infty(\mathbb{R}^+)} + \Delta t_n F_1(0) \text{ which completes the proof.}$$

Proposition 2.4. *Under the hypotheses of proposition 2.1, the function v_h^n satisfies:*

$$\|v_h^n\|_{L^1(\mathbb{R}^+)} \leq t_n F_1(0)$$

for $n \geq 0$.

Proof: From (2.16), we get if $\lambda > 0$

$$\sum_{i=1}^N h_i v_i^{n+\frac{1}{2}} = \sum_{i=1}^N h_i v_i^n \left(1 - \lambda \frac{\Delta t_n}{h_i} x_i^2 v_i^n \right) + \lambda \Delta t_n \sum_{i=1}^N x_{i-1}^2 (v_{i-1}^n)^2,$$

hence $\left\|v_h^{n+\frac{1}{2}}\right\|_{L^1(\mathbb{R}^+)} \leq \|v_h^n\|_{L^1(\mathbb{R}^+)}.$

For $\lambda < 0$, we obtain the same inequality.

Besides from (2.12), we get:

$$\|v_h^{n+1}\|_{L^1(\mathbb{R}^+)} \leq \left\|v_h^{n+\frac{1}{2}}\right\|_{L^1(\mathbb{R}^+)} - \Delta t_n F_{1N} + \Delta t_n F_{10}.$$

and we deduce the result.

We prove now that under some hypothesis on the sequence (h_i) , the function xv_h^n is bounded in $L^\infty(\mathbb{R}^+)$ and it is possible to choose the nodes $(x_i)_{1 \leq i \leq N}$ such that the stability condition is not too restrictive.

We define the function $\hat{v}_h^n$ by:

$$\hat{v}_h^n|_{I_i} = \hat{v}_i^n = x_i v_i^n, \quad 1 \leq i \leq N, \quad \hat{v}_h^n|_{I_{N+1}} = \hat{v}_{N+1}^n = 0.$$

Proposition 2.5. *If the following stability condition*

$$(2.18) \quad \sup_{i \geq 1} \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i-1}^n) \leq 1 \text{ if } \lambda > 0 \text{ and } \sup_{i \geq 1} |\lambda| \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i+1}^n) \leq 1 \text{ if } \lambda < 0$$

is satisfied and if the sequence $(h_i)_{1 \leq i \leq N}$ satisfy: There exists a positive constant c such that

$$(2.19) \quad \frac{x_i}{x_{i+1}} \frac{h_{i+1}}{h_i + h_{i+1}} - \frac{x_{i-1}}{x_i} \frac{h_i}{h_i + h_{i-1}} \geq -c \frac{h_i}{x_i}, \quad 1 \leq i \leq N-1$$

and

$$(2.20) \quad \frac{x_N}{h_N + 2(1 + x_N)} - \frac{x_{N-1}}{x_N} \frac{h_N}{h_N + h_{N-1}} \geq -c \frac{h_N}{x_N},$$

then if $\Delta t \leq \Delta t_0$, Δt_0 depending on c, c_1, c_2 , the following estimate holds:

$$(2.21) \quad \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)} \leq e^{Ct_n} \|\hat{F}\|_{L^\infty(\mathbb{R}^+)}$$

for $n \geq 0$ and C is a constant depending on c, c_1, c_2 .

Proof: We get immediately from (2.10)

$$\hat{v}_i^{n+\frac{1}{2}} = \hat{v}_i^n - \lambda \frac{\Delta t_n}{h_i} x_i ((\hat{v}_i^n)^2 - (\hat{v}_{i-1}^n)^2) \text{ if } \lambda > 0$$

which may be written:

$$\hat{v}_i^{n+\frac{1}{2}} = \hat{v}_i^n \left(1 - \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i-1}^n) \right) + \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i-1}^n) \hat{v}_{i-1}^n$$

and if (2.18) is satisfied, we obtain:

$$(2.22) \quad \left\|\hat{v}_h^{n+\frac{1}{2}}\right\|_{L^\infty(\mathbb{R}^+)} \leq \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)}.$$

If $\lambda < 0$, we get from (2.11)

$$\hat{v}_i^{n+\frac{1}{2}} = \hat{v}_i^n \left(1 + \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_{i+1}^n + \hat{v}_i^n) \right) - \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_{i+1}^n + \hat{v}_i^n) \hat{v}_{i+1}^n$$

and the estimate (2.22) holds if (2.18) is satisfied.

Further the following equality results of (2.12) for $1 \leq i \leq N$

$$\begin{aligned} (2.23) \quad & \hat{v}_i^{n+1} + \frac{\Delta t_n}{h_i} (\alpha_i + \alpha_{i-1} - \delta_i \gamma_i + (1 - \delta_{i-1}) \gamma_{i-1}) \hat{v}_i^{n+1} \\ & - \frac{\Delta t_n}{h_i} (\alpha_{i-1} - \delta_{i-1} \gamma_{i-1}) \frac{x_i}{x_{i-1}} \hat{v}_{i-1}^{n+1} - \frac{\Delta t_n}{h_i} (\alpha_i + (1 - \delta_i) \gamma_i) \frac{x_i}{x_{i+1}} \hat{v}_{i+1}^{n+1} \\ & = \hat{v}_i^{n+\frac{1}{2}} - \frac{\Delta t_n}{h_i} x_i (F_{1i} - F_{1,i-1}) \end{aligned}$$

which may be written

$$(I + \Delta t_n C_h) \hat{V}_h^{n+1} = \hat{V}_h^{n+\frac{1}{2}} - \Delta t_n \hat{F}_h$$

where $\hat{F}_h$ is the vector of components : $\hat{F}_i = x_i \frac{F_{1i} - F_{1,i-1}}{h_i}$, $1 \leq i \leq N$. C_h is a tridiagonal matrix ($N \times N$).

Besides, we have $c_{ii} > 0$, $c_{ij} \leq 0$ for $i \neq j$, $1 \leq i \leq N$.

We prove that if (2.19) and (2.20) are satisfied, there exists a positive constant $\hat{c}$ depending on

c_1 , c_2 , c such that: $\sum_{j=1}^N c_{ij} \geq -\hat{c}$, $1 \leq i \leq N$.

We deduce from (2.2):

$$\begin{aligned} \sum_{j=1}^N c_{ij} &= \frac{1}{h_i} \left(\alpha_i \frac{h_{i+1}}{x_{i+1}} - \alpha_{i-1} \frac{h_i}{x_{i-1}} - \gamma_i + \gamma_{i-1} + \gamma_i (1 - \delta_i) \frac{h_{i+1}}{x_{i+1}} + \gamma_{i-1} \delta_{i-1} \frac{h_i}{x_{i-1}} \right), \quad 1 \leq i \leq N-1 \\ \sum_{j=1}^N c_{Nj} &= \frac{1}{h_N} \left(\alpha_N - \alpha_{N-1} \frac{h_N}{x_{N-1}} - \gamma_N + \gamma_{N-1} \frac{x_N}{x_{N-1}} \right). \end{aligned}$$

Let us denote $A_i^1 = \frac{1}{h_i} \left(\alpha_i \frac{h_{i+1}}{x_{i+1}} - \alpha_{i-1} \frac{h_i}{x_{i-1}} \right) = \frac{k^2 x_i}{h_i} \left(\frac{x_i}{x_{i+1}} \frac{h_{i+1}}{h_i + h_{i+1}} - \frac{x_{i-1}}{x_i} \frac{h_i}{h_{i-1} + h_i} \right)$,

$1 \leq i \leq N-1$,

$$A_N^1 = \frac{1}{h_N} \left(\alpha_N - \alpha_{N-1} \frac{h_N}{x_{N-1}} \right) = \frac{k^2 x_N}{h_N} \left(\frac{x_N}{h_N + 2(1 + x_N)} - \frac{x_{N-1}}{x_N} \frac{h_N}{h_N + h_{N-1}} \right);$$

and $A_i^2 = \frac{1}{h_i} \left(-\gamma_i + \gamma_{i-1} + \gamma_i (1 - \delta_i) \frac{h_{i+1}}{x_{i+1}} + \gamma_{i-1} \delta_{i-1} \frac{h_i}{x_{i-1}} \right)$, $2 \leq i \leq N$,

$$A_1^2 = \frac{1}{h} \left(-\delta_1 - \frac{1 - \delta_1}{2} \right) \gamma_1.$$

We get $\sum_{j=1}^N c_{ij} = A_i^1 + A_i^2$.

From (2.19) and (2.20), it follows that $A_i^1 \geq -ck^2$, $1 \leq i \leq N$.

Further, we have : $A_1^2 = -\phi(x_1)(\delta_1 + \frac{1 - \delta_1}{2})$

and for $i \geq 2$, $A_i^2 = -\frac{g_1(x_i) - g_1(x_{i-1})}{x_i - x_{i-1}} + (1 - \delta_i) \frac{x_i h_{i+1}}{x_{i+1} h_i} \phi(x_i) + \delta_{i-1} \phi(x_{i-1})$,

We deduce from (2.2) $A_i^2 \geq -(c_1 + c_2)$ for $x_i \leq \sigma_2$.

For $x \geq \sigma_2$, the function g_1 is negative, so $\delta_i = 1$, $A_i^2 = \frac{-x_i(\phi(x_i) - \phi(x_{i-1}))}{h_i}$
and $A_i^2 \geq 0$, since ϕ is decreasing.

Finally, we obtain $\sum_{j=1}^N c_{ij} \geq -\hat{c}$, $1 \leq i \leq N$, with $\hat{c} = ck^2 + c_1 + c_2$.

and $\|\hat{v}_h^{n+1}\|_{L^\infty(\mathbb{R}^+)} \leq \frac{1}{1 - \hat{c}\Delta t_n} \left(\|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)} + \Delta t_n \|\hat{F}\|_{L^\infty(\mathbb{R}^+)} \right)$.

The estimate (2.21) follows.

Let us define now a sequence (x_i) , satisfying (2.19) and (2.20):

We set: $x_i = ih$, $0 \leq i \leq n_0$ with $n_0 h = 1$,
then $x_i = e^{\theta h} x_{i-1}$ for $i \geq n_0 + 1$ and $\theta \geq 1$

We get: $\frac{x_i}{x_{i+1}} \frac{h_{i+1}}{h_{i+1} + h_i} - \frac{x_{i-1}}{x_i} \frac{h_i}{h_i + h_{i-1}} > 0$, $i \leq n_0 + 1$,

$\frac{x_i}{x_{i+1}} \frac{h_{i+1}}{h_{i+1} + h_i} - \frac{x_{i-1}}{x_i} \frac{h_i}{h_i + h_{i-1}} = 0$, $i > n_0 + 1$

$\frac{x_N}{h_N + 2(1 + x_N)} - \frac{x_{N-1}}{x_N} \frac{h_N}{h_N + h_{N-1}} \geq -\frac{1}{2x_N}$; then (2.20) will be satisfied if $h_N \geq \frac{\epsilon}{2}$, that is
 $N = \mathcal{O}\left(\frac{|\ln h|}{h}\right)$ or $x_N = \mathcal{O}\left(\frac{1}{h}\right)$.

Besides we have $\frac{x_i}{h_i} = \mathcal{O}\left(\frac{1}{h}\right)$ $1 \leq i \leq N$, and the stability condition may be written $\frac{\Delta t_n}{h} \leq C$,
that is the classical stability condition for hyperbolic problems.

Proposition 2.6. *Under the hypotheses of proposition 2.5, there exists a positive constant C depending on T, f, g_1 such that for $t_n \leq T$, the following estimate holds:*

$$\|v_h^n\|_{L^\infty(\mathbb{R}^+)} \leq C.$$

Proof: For $\lambda > 0$, we get from (2.10):

$$v_i^{n+\frac{1}{2}} = v_i^n \left(1 - \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i-1}^n) \right) + \lambda \frac{\Delta t_n}{h_i} x_{i-1} (\hat{v}_i^n + \hat{v}_{i-1}^n) v_{i-1}^n$$

and by using (2.18), we obtain: $\|v_h^{n+\frac{1}{2}}\|_{L^\infty(\mathbb{R}^+)} \leq \|v_h^n\|_{L^\infty(\mathbb{R}^+)}.$

For $\lambda < 0$, we get:

$$v_i^{n+\frac{1}{2}} = v_i^n \left(1 + \lambda \frac{\Delta t_n}{h_i} x_i (\hat{v}_i^n + \hat{v}_{i+1}^n) \right) - \lambda \frac{\Delta t_n}{h_i} x_{i+1} (\hat{v}_i^n + \hat{v}_{i+1}^n) v_{i+1}^n$$

and by using (2.18), we obtain:

$$\|v_h^{n+\frac{1}{2}}\|_{L^\infty(\mathbb{R}^+)} \leq \|v_h^n\|_{L^\infty(\mathbb{R}^+)} \left(1 + 2|\lambda| \Delta t_n \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)} \right).$$

Besides, it follows from (2.13) that

$$(1 - c_1 \Delta t_n) \|v_h^{n+1}\|_{L^\infty(\mathbb{R}^+)} \leq \|v_h^{n+\frac{1}{2}}\|_{L^\infty(\mathbb{R}^+)} + \Delta t_n \|F'_1\|_{L^\infty(\mathbb{R}^+)}.$$

This concludes the proof.

Proposition 2.7. *Under the hypotheses of proposition 2.5, there a positive constant C depending on T, f, g_1 such that for $t_n \leq T$, the following estimate holds:*

$$\text{Var}(v_h^n; \mathbb{R}^+) \leq C.$$

Proof: For $\lambda > 0$, we have the equality:

$$v_{i+1}^{n+\frac{1}{2}} - v_i^{n+\frac{1}{2}} = v_{i+1}^n - v_i^n - \lambda \frac{\Delta t_n}{h_{i+1}} (x_{i+1}^2 (v_{i+1}^n)^2 - x_i^2 (v_i^n)^2) + \lambda \frac{\Delta t_n}{h_i} (x_i^2 (v_i^n)^2 - x_{i-1}^2 (v_{i-1}^n)^2)$$

for $1 \leq i \leq N-1$, and

$$v_{N+1}^{n+\frac{1}{2}} - v_N^{n+\frac{1}{2}} = v_{N+1}^n - v_N^n + \lambda \frac{\Delta t_n}{h_N} (x_N^2 (v_N^n)^2 - x_{N-1}^2 (v_{N-1}^n)^2).$$

It follows that:

$$\begin{aligned} v_{i+1}^{n+\frac{1}{2}} - v_i^{n+\frac{1}{2}} &= (v_{i+1}^n - v_i^n)(1 - \mu_i^n) + \mu_{i-1}^n (v_i^n - v_{i-1}^n) \\ &\quad - 2\lambda \Delta t_n x_i (v_{i+1}^n + v_i^n)(v_{i+1}^n - v_i^n) - \lambda \Delta t_n h_{i+1} (v_{i+1}^n)^2 - \lambda \Delta t_n h_i (v_i^n)^2 \end{aligned}$$

$$\text{with } \mu_i^n = \lambda \frac{\Delta t_n}{h_{i+1}} x_i^2 (v_{i+1}^n + v_i^n), \quad 1 \leq i \leq N-1, \quad \mu_N^n = 0.$$

From the stability condition (2.18), we get $0 \leq \mu_i^n \leq 1$ and we deduce

$$\text{Var}(v_h^{n+\frac{1}{2}}; \mathbb{R}^+) \leq \text{Var}(v_h^n; \mathbb{R}^+) (1 + 4\lambda \Delta t_n \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)}) + 2\lambda \Delta t_n \|v_h^n\|_{L^\infty(\mathbb{R}^+)} \|v_h^n\|_{L^1(\mathbb{R}^+)}.$$

We have an analogous estimate for $\lambda < 0$.

Furthermore, from (2.12), we obtain the following equalities:

$$\begin{aligned}
& v_{i+1}^{n+1} - v_i^{n+1} + \Delta t_n \left(\frac{\alpha_i + (1 - \delta_i)\gamma_i}{h_i} + \frac{\alpha_i - \delta_i\gamma_i}{h_{i+1}} - \frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} \right) (v_{i+1}^{n+1} - v_i^{n+1}) \\
& - \Delta t_n \frac{\alpha_{i+1} + (1 - \delta_{i+1})\gamma_{i+1}}{h_{i+1}} (v_{i+2}^{n+1} - v_{i+1}^{n+1}) - \Delta t_n \frac{\alpha_{i-1} - \delta_{i-1}\gamma_{i-1}}{h_i} (v_i^{n+1} - v_{i-1}^{n+1}) \\
& = v_{i+1}^{n+\frac{1}{2}} - v_i^{n+\frac{1}{2}} + \Delta t_n \left(\frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} - \frac{\gamma_i - \gamma_{i-1}}{h_i} \right) v_i^{n+1} - \Delta t_n \left(\frac{F_{1,i+1} - F_{1i}}{h_{i+1}} - \frac{F_{1i} - F_{1,i-1}}{h_i} \right),
\end{aligned}$$

for $1 \leq i \leq N-1$.

$$\begin{aligned}
& v_{N+1}^{n+1} - v_N^{n+1} + \Delta t_n \left(\frac{\alpha_N}{h_N} (v_{N+1}^{n+1} - v_N^{n+1}) - \frac{\alpha_{N-1} - \gamma_{N-1}}{h_N} (v_N^{n+1} - v_{N-1}^{n+1}) \right) \\
& = v_{N+1}^{n+\frac{1}{2}} - v_N^{n+\frac{1}{2}} - \Delta t_n \frac{\gamma_N - \gamma_{N-1}}{h_N} + \Delta t_n \frac{F_{1N} - F_{1N-1}}{h_N}.
\end{aligned}$$

It follows that:

$$\begin{aligned}
& \sum_{i=1}^{N-1} \left(1 - \Delta t_n \frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} \right) |v_{i+1}^{n+1} - v_i^{n+1}| + v_N^{n+1} \leq Var(v_h^{n+\frac{1}{2}}; \mathbb{R}^+) \\
& + \Delta t_n \sum_{i=1}^{N-1} \left| \frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} - \frac{\gamma_i - \gamma_{i-1}}{h_i} \right| v_i^{n+1} \\
& + \Delta t_n \sum_{i=1}^{N-1} \left| \frac{F_{1i} - F_{1,i-1}}{h_{i+1}} - \frac{F_{1i} - F_{1,i-1}}{h_i} \right| + \Delta t_n \left| \frac{F_{1N} - F_{1,N-1}}{h_N} \right|.
\end{aligned}$$

It results from (2.2): $1 - \Delta t_n \frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} \geq 1 - c_1 \Delta t_n$.

Besides, we have: $\sum_{i=1}^{N-1} \left| \frac{\gamma_{i+1} - \gamma_i}{h_{i+1}} - \frac{\gamma_i - \gamma_{i-1}}{h_i} \right| v_i^{n+1} \leq Var(g'_1; \mathbb{R}^+) \|v_h^{n+1}\|_{L^\infty(\mathbb{R}^+)}$

and $\sum_{i=1}^{N-1} \left| \frac{F_{1,i+1} - F_{1i}}{h_{i+1}} - \frac{F_{1i} - F_{1,i-1}}{h_i} \right| \leq Var(F'_1; \mathbb{R}^+)$.

It follows that:

$$(1 - c_1 \Delta t_n) \text{Var}(v_h^{n+1}; \mathbb{R}^+) \leq \text{Var}(v_h^{n+\frac{1}{2}}; \mathbb{R}^+) + C \Delta t_n \|v_h^{n+1}\|_{L^1(\mathbb{R}^+)} + \Delta t_n \text{Var}(F'_1, \mathbb{R}^+)$$

and with (2.2), we get:

$$(1 - c_1 \Delta t_n) \text{Var}(v_h^{n+1}; \mathbb{R}^+) \leq \text{Var}(v_h^n; \mathbb{R}^+) (1 + 4\lambda \Delta t_n \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)})$$

$$+ 2\lambda \Delta t_n \|v_h^n\|_{L^\infty(\mathbb{R}^+)} \|v_h^n\|_{L^1(\mathbb{R}^+)} + C \Delta t_n \|v_h^{n+1}\|_{L^1(\mathbb{R}^+)} + \Delta t_n \text{Var}(F'_1; \mathbb{R}^+).$$

This concludes the proof.

2.3. Convergence of the scheme and uniqueness of the solution.

From all these estimates, we can deduce the convergence of the numerical solution to a weak solution and we prove that this solution is unique.

A function u_1 is called a *weak solution* of (2.1) if $u_1 \in C([0, T]); W^{1,\infty}(\mathbb{R}^+)$, $x \frac{\partial u_1}{\partial x} \in C(0, T; L^\infty(\mathbb{R}^+))$, and

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^+} u_1 \frac{\partial \phi}{\partial t} dx dt - \int_{\mathbb{R}^+} u_1(T) \phi(T) dx - \frac{k^2}{2} \int_0^T \int_{\mathbb{R}^+} \frac{\partial u_1}{\partial x} \frac{\partial}{\partial x} (x^2 \phi) dx dx \\ & + \int_0^T \int_{\mathbb{R}^+} g_1 \frac{\partial u_1}{\partial x} \phi dx dt - \lambda \int_0^T \int_{\mathbb{R}^+} x^2 \left(\frac{\partial u_1}{\partial x} \right)^2 \phi dx dt = \int_0^T \int_{\mathbb{R}^+} F_1 \phi dx dt \end{aligned}$$

for any function ϕ with a compact support in $[0, T] \times \mathbb{R}^+$, $\phi \in C^1(0, T \times \mathbb{R}^+)$.

Theorem 2.8. *Problem (2.1) admits at most one weak solution.*

Proof: Let u_1 and $\hat{u}_1$ two weak solutions of (2.1). We denote $w = u_1 - \hat{u}_1$. The function w satisfies:

$$(2.24) \quad \frac{\partial w}{\partial t} - \frac{1}{2} k^2 x^2 \frac{\partial^2 w}{\partial x^2} - g_1(x) \frac{\partial w}{\partial x} + \lambda x^2 \frac{\partial w}{\partial x} \left(\frac{\partial u_1}{\partial x} + \frac{\partial \hat{u}_1}{\partial x} \right) = 0.$$

Let us denote by ψ a function in $C^1(\mathbb{R}^+)$ satisfying:

$0 \leq \psi(x) \leq 1$, $\psi(x) = 1$ if $0 \leq x \leq 1$, $\psi(x) = 0$ if $x \geq 2$ and ψ decreasing on $(1, 2)$ and we define $\psi_\nu(x) = \psi(\frac{x}{\nu})$, $x \geq 0$, $\nu > 0$.

By multiplying (2.24) by $\psi_\nu \frac{w}{(1+x)^2}$, and integrating on $\mathbb{R}^+$, we get:

$$\frac{1}{2} \frac{d}{dt} \left(\int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx \right) + \frac{1}{2} k^2 \int_0^{+\infty} \left(\frac{\partial w}{\partial x} \right)^2 \frac{x^2}{(1+x)^2} \psi_\nu dx + \frac{1}{2} \int_0^{+\infty} w^2 \frac{g_1(x)}{(1+x)^2} \frac{d\psi_\nu}{dx} dx$$

$$\begin{aligned}
&= -\frac{k^2}{2} \int_0^{+\infty} \frac{\partial w}{\partial x} w \frac{x^2}{(1+x)^2} \frac{d\psi_\nu}{dx} dx - k^2 \int_0^{+\infty} \frac{\partial w}{\partial x} w \frac{x}{(1+x)^3} \psi_\nu dx \\
(2.25) \quad &- \frac{1}{2} \int_0^{+\infty} w^2 \psi_\nu \frac{d}{dx} \left(\frac{g_1(x)}{(1+x)^2} \right) dx - \lambda \int_0^{+\infty} \frac{\partial w}{\partial x} w \left(\frac{\partial u_1}{\partial x} + \frac{\partial \hat{u}_1}{\partial x} \right) \frac{x^2}{(1+x)^2} \psi_\nu dx.
\end{aligned}$$

We estimate now each term of this equality.

We have: $\int_0^{+\infty} w^2 \frac{g_1(x)}{(1+x)^2} \frac{d\psi_\nu}{dx} dx \geq 0$ if $\nu \geq \sigma_2$ since $g_1(x) \leq 0$ and $\psi'_\nu(x) \leq 0$.

Besides, we have: $\psi'_\nu(x) = \frac{1}{\nu} \psi'(\frac{x}{\nu})$ and since $x \frac{\partial w}{\partial x} \in C(0, T; L^\infty(\mathbb{R}^+))$, $w \in C(0, T; L^\infty(\mathbb{R}^+))$, we obtain:

$$\frac{k^2}{2} \left| \int_0^{+\infty} \frac{\partial w}{\partial x} w \frac{x^2}{(1+x)^2} \frac{d\psi_\nu}{dx} dx \right| \leq \frac{C}{\nu} \quad (C \text{ depending on } \|x \frac{\partial w}{\partial x}\|_{L^\infty(\mathbb{R}^+)} \text{ and } \|w\|_{L^\infty(\mathbb{R}^+)}).$$

We estimate the second term of the second member of (2.25) and we get:

$$\begin{aligned}
k^2 \left| \int_0^{+\infty} \frac{\partial w}{\partial x} w \frac{x}{(1+x)^3} \psi_\nu dx \right| &\leq k^2 \left(\int_0^{+\infty} \frac{w^2}{(1+x)^2} \psi_\nu dx \right)^{\frac{1}{2}} \left(\int_0^{+\infty} \left(\frac{\partial w}{\partial x} \right)^2 \frac{x^2}{(1+x)^4} \psi_\nu dx \right)^{\frac{1}{2}} \\
&\leq \alpha k^2 \int_0^{+\infty} \left(\frac{\partial w}{\partial x} \right)^2 \frac{x^2}{(1+x)^2} \psi_\nu dx + \frac{k^2}{4\alpha} \int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx,
\end{aligned}$$

$\alpha > 0$.

$$\text{Further, we have: } \left| \int_0^{+\infty} w^2 \psi_\nu \frac{d}{dx} \left(\frac{g_1(x)}{(1+x)^2} \right) dx \right| \leq \int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} \left| \frac{g'_1(x)(1+x) - 2g_1(x)}{1+x} \right| dx.$$

From the hypotheses on the function g_1 , we get

$$\left| \frac{g'_1(x)(1+x) - 2g_1(x)}{1+x} \right| \leq \left| \frac{\phi(x) + x\phi'(x)}{1+x} \right| + |\phi(x) - x\phi'(x)|$$

and this quantity is bounded. Then, we obtain:

$$\left| \int_0^{+\infty} w^2 \psi_\nu \frac{d}{dx} \left(\frac{g_1(x)}{(1+x)^2} \right) dx \right| \leq C \int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx.$$

It remains to study the last term of (2.25):

Since $x \frac{\partial u_1}{\partial x}$ and $x \frac{\partial \hat{u}_1}{\partial x} \in C(0, T; L^\infty(\mathbb{R}^+))$, we get:

$$\left| \int_0^{+\infty} \frac{\partial w}{\partial x} w \left(\frac{\partial u_1}{\partial x} + \frac{\partial \hat{u}_1}{\partial x} \right) \frac{x^2}{(1+x)^2} \psi_\nu dx \right| \leq C \int_0^{+\infty} \left| \frac{\partial w}{\partial x} \right| |w| \frac{x}{(1+x)^2} \psi_\nu dx$$

$$\leq \alpha k^2 \int_0^{+\infty} \left(\frac{\partial w}{\partial x} \right)^2 \frac{x^2}{(1+x)^2} \psi_\nu dx + \frac{C^2}{4\alpha k^2} \int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx.$$

We deduce from all these estimates:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx \right) + k^2 \left(\frac{1}{2} - 2\alpha \right) \int_0^{+\infty} \left(\frac{\partial w}{\partial x} \right)^2 \frac{x^2}{(1+x)^2} \psi_\nu dx \\ \leq \frac{C}{\nu} + C_1 \int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx. \end{aligned}$$

If we choose $\alpha < \frac{1}{4}$, we obtain by using the Gronwall's lemma:

$$\int_0^{+\infty} w^2 \frac{\psi_\nu}{(1+x)^2} dx \leq \frac{2CT}{\nu} e^{2C_1 t}, \quad 0 \leq t \leq T$$

and if $\nu \rightarrow +\infty$, we get $\int_0^{+\infty} \frac{w^2}{(1+x)^2} dx = 0$

and we deduce $w = 0$ and the problem admits at most one solution.

We prove now the convergence of the numerical solution to this weak solution and thus, we obtain the existence of a solution.

We define the functions $u_{1h\Delta t}$ and $v_{1h\Delta t}$ by

$$\begin{aligned} u_{1h\Delta t} &= u_h^n + \frac{t - t_n}{\Delta t_n} (u_{1h}^{n+1} - u_{1h}^n), \\ v_{1h\Delta t} &= v_h^n + \frac{t - t_n}{\Delta t_n} (v_h^{n+1} - v_h^n), \quad t_n \leq t \leq t_{n+1}. \end{aligned}$$

Theorem 2.9. *Assume that the hypotheses of proposition 2.5 are satisfied. The sequence $u_{1h\Delta t}$ converges uniformly to the weak solution of (2.1) on any compact of $[0, T] \times \mathbb{R}^+$.*

Proof: The functions $(u_{1h\Delta t})$ are uniformly bounded in $C(0, T; W^{1,\infty}(\mathbb{R}^+))$.

Further, for $R > 0$, we get the estimate:

$$\begin{aligned} \left\| \frac{u_{1h}^{n+1} - u_{1h}^n}{\Delta t_n} \right\|_{L^1(0, R)} &\leq \frac{k^2 R^2}{2} \text{Var}(v_h^{n+1}; \mathbb{R}^+) + C(g_1, R) \|v_h^{n+1}\|_{L^1(\mathbb{R}^+)} \\ &+ |\lambda| R \|\hat{v}_h^n\|_{L^\infty(\mathbb{R}^+)} \|v_h^n\|_{L^1(\mathbb{R}^+)} + \|F_1\|_{L^1(\mathbb{R}^+)} \end{aligned}$$

with $C(g_1, R) = \sup_{x \leq R} |g_1(x)|$.

Thus, the time derivatives of $u_{h\Delta t}$ are uniformly bounded in $L^\infty(0, T; L^1(0, R))$ for any $R > 0$.

So, we can extract from the sequence $(u_{h\Delta t})$ a subsequence, again labeled $u_{h\Delta t}$ which converges uniformly on any compact subset of $[0, T] \times \mathbb{R}^+$ to a function u_1 [7].

The functions $v_{h\Delta t}$ are uniformly bounded in $C(0, T; BV(\mathbb{R}^+))$; besides, we have:

(2.26)

$$\|v_h^{n+1} - v_h^n\|_{H^{-2}(0, R)} = \sup_{\phi \in H_0^2(0, R)} \frac{\langle v_h^{n+1} - v_h^n, \phi \rangle}{\|\phi\|_{H_0^2(0, R)}} = \sup_{\phi \in H_0^2(0, R)} \frac{\langle u_h^{n+1} - u_h^n, \phi_x \rangle}{\|\phi\|_{H_0^2(0, R)}}$$

$$\text{and } \left\| \frac{v_h^{n+1} - v_h^n}{\Delta t_n} \right\|_{H^{-2}(0, R)} \leq C \left\| \frac{u_h^{n+1} - u_h^n}{\Delta t_n} \right\|_{L^1(0, R)} \leq C(R).$$

Then, we can extract from $(v_{h\Delta t})$ a subsequence, again labeled $v_{h\Delta t}$ which converges to a function $v = \frac{\partial u_1}{\partial x}$ in $C(0, T; L^1(Q))$ for any compact $Q \subset \mathbb{R}^+$ [7].

We get easily that u_1 is a weak solution. Since this solution is unique, all the sequence is converging to u_1 .

So, we have obtained the following result:

Theorem 2.10. *Problem (2.1) admits a unique weak solution.*

The following figure represents $a(0)$ with a time maturity [3] equal to 1 for different values of ρ . the values of the other parameters are those proposed in [1]: $k = 0.4$, $\delta = 2$, $\sigma_1 = 0.153$, $\mu = 0.7$, $\sigma_0 = 0.01$.




FIGURE 1. a with $T=1$

3. COMPUTATION OF u_2

u_2 is solution of

$$\frac{\partial u_2}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 u_2}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 u_2}{\partial y^2} - \rho k x^2 y \frac{\partial^2 u_2}{\partial x \partial y} - g_2(x) \frac{\partial u_2}{\partial x}$$

$$-(1 - \rho^2) k^2 x^2 \frac{\partial u_1}{\partial x} \frac{\partial u_2}{\partial x} = 0.$$

The initial condition is the payoff function. If the European option is a put, the payoff function is given by: $F(y) = \text{Max}(E - y, 0)$ if E is the exercise price.

3.1. Existence and uniqueness of the solution.

In order to obtain a bilinear form satisfying Garding's inequality, we make a change of unknown.

We denote: $\hat{u}_2 = e^{-\alpha x} u_2$, $\alpha > 0$.

The function $\hat{u}_2$ is solution of:

$$\begin{aligned} & \frac{\partial \hat{u}_2}{\partial t} - \frac{k^2}{2} x^2 \frac{\partial^2 \hat{u}_2}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 \hat{u}_2}{\partial y^2} - \rho k x^2 y \frac{\partial^2 \hat{u}_2}{\partial x \partial y} - \frac{\partial \hat{u}_2}{\partial x} \left(\alpha k^2 x^2 + g_2(x) + (1 - \rho^2) k^2 x^2 \frac{\partial u_1}{\partial x} \right) \\ & - \rho \alpha k x^2 y \frac{\partial \hat{u}_2}{\partial y} - \left(\frac{\alpha^2 k^2}{2} x^2 + \alpha g_2(x) + (1 - \rho^2) \alpha k^2 x^2 \frac{\partial u_1}{\partial x} \right) \hat{u}_2 = 0, \end{aligned}$$

with the initial condition: $\hat{u}_2(0) = e^{-\alpha x} F(y)$.

We define a variational formulation of this problem.

Let us consider the following space $\hat{V}_2$ defined by:

$$\hat{V}_2 = \{ v \in \mathcal{D}'(\Omega) / v \in L^2(\Omega), xv \in L^2(\Omega), xv_x \in L^2(\Omega), xyv_y \in L^2(\Omega) \}$$

with $\Omega = \mathbb{R}^+ \times \mathbb{R}^+$.

This space with the norm:

$$\|v\| = \left(\int_{\Omega} v^2 + x^2 v^2 + x^2 \left(\frac{\partial v}{\partial x} \right)^2 + x^2 y^2 \left(\frac{\partial v}{\partial y} \right)^2 \right)^{\frac{1}{2}}$$

is a Hilbert space and $\mathcal{D}(\Omega)$ is dense in $\hat{V}_2$ [3].

Besides, we have the estimates:

$$\|xv\|_{L^2(\Omega)} \leq 2 \left\| xy \frac{\partial v}{\partial y} \right\|_{L^2(\Omega)}, \|v\|_{L^2(\Omega)} \leq 2 \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}$$

and the semi-norm:

$$\|v\|_2 = \left(\left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 + \left\| xy \frac{\partial v}{\partial y} \right\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}}$$

is a norm in $\hat{V}_2$ equivalent to the norm $\|\cdot\|_{\hat{V}_2}$ [3].

We define on $\hat{V}_2 \times \hat{V}_2$ the bilinear form $\hat{b}$ by :

$$\begin{aligned} \forall u, v \in \hat{V}_2, \hat{b}(u, v) &= \frac{k^2}{2} \int_{\Omega} \frac{\partial u}{\partial x} \frac{\partial}{\partial x} (x^2 v) dx dy + \frac{1}{2} \int_{\Omega} x^2 \frac{\partial u}{\partial y} \frac{\partial}{\partial y} (y^2 v) dx dy \\ &+ \rho k \int_{\Omega} x^2 \frac{\partial u}{\partial x} \frac{\partial}{\partial y} (y v) dx dy - \int_{\Omega} (\alpha k^2 x^2 + g_2(x) + (1 - \rho^2) k^2 x^2 \frac{\partial u_1}{\partial x}) \frac{\partial u}{\partial x} v dx dy \\ &- \rho \alpha k \int_{\Omega} x^2 y \frac{\partial u}{\partial y} v dx dy - \int_{\Omega} \left(\frac{\alpha^2 k^2}{2} x^2 + \alpha g_2(x) + (1 - \rho^2) \alpha k^2 x^2 \frac{\partial u_1}{\partial x} \right) u v dx dy. \end{aligned}$$

We have proved in the preceding section that the function $x \frac{\partial u_1}{\partial x}$ is in $C(0, T; L^\infty(\mathbb{R}^+))$.

Besides, we have $g_2(x) = -\delta x(x - \sigma_1) - \rho k x f(x)$.

We denote $f_1(x) = x f(x)$ and we assume that $f_1, f'_1 \in L^\infty(\mathbb{R}^+)$.

It is clear that the bilinear form $\hat{b}$ is continue on $\hat{V}_2 \times \hat{V}_2$.

We prove now that it is possible to choose $\alpha > 0$ such that this bilinear form satisfies Garding's inequality.

Proposition 3.1. *If $|\rho| < 1$ and $k < \frac{2}{3}\delta$, there exists positive constants C, c, α depending on $\delta, \sigma_1, k, \rho$ such that*

$$\forall v \in \hat{V}_2, \hat{b}(v, v) \geq C \|v\|_2^2 - c \|v\|_{L^2(\Omega)}^2.$$

Proof: We have the equality:

$$\hat{b}(v, v) = \sum_{i=1}^6 B_i$$

with

$$\begin{aligned} B_1 &= \frac{k^2}{2} \int_{\Omega} \frac{\partial v}{\partial x} \frac{\partial}{\partial x} (x^2 v) dx dy; \quad B_2 = \frac{1}{2} \int_{\Omega} x^2 \frac{\partial v}{\partial y} \frac{\partial}{\partial y} (y^2 v) dx dy; \quad B_3 = \rho k \int_{\Omega} x^2 \frac{\partial v}{\partial x} \frac{\partial}{\partial y} (y v) dx dy; \\ B_4 &= - \int_{\Omega} \left(\alpha k^2 x^2 + g_2(x) \right) + (1 - \rho^2) k^2 x^2 \frac{\partial u_1}{\partial x} \Big) v \frac{\partial v}{\partial x} dx dy; \quad B_5 = - \rho \alpha k \int_{\Omega} x^2 y \frac{\partial v}{\partial y} v dx dy; \\ B_6 &= - \int_{\Omega} \left(\frac{\alpha^2 k^2}{2} x^2 + \alpha g_2(x) + (1 - \rho^2) \alpha k^2 x^2 \frac{\partial u_1}{\partial x} \right) v^2 dx dy. \end{aligned}$$

It is easily seen, for $\epsilon_i > 0, 1 \leq i \leq 4$

$$B_1 = \frac{k^2}{2} \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 - \frac{k^2}{2} \|v\|_{L^2(\Omega)}^2,$$

$$B_2 = \frac{1}{2} \left\| xy \frac{\partial v}{\partial y} \right\|_{L^2(\Omega)}^2 - \frac{1}{2} \|xv\|_{L^2(\Omega)}^2,$$

$$B_3 = \rho k \int_{\Omega} (x^2 y \frac{\partial v}{\partial x} \frac{\partial v}{\partial y} - xv^2) dx dy \text{ and this term is bounded from below:}$$

$$|B_3| \geq -\frac{|\rho|}{2} \left(\left\| xy \frac{\partial v}{\partial y} \right\|_{L^2(\Omega)}^2 + k^2 \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 \right) - |\rho| \left(\epsilon_1 \|xv\|_{L^2(\Omega)}^2 + \frac{k^2}{4\epsilon_1} \|v\|_{L^2(\Omega)}^2 \right), \quad \epsilon_1 > 0,$$

$$B_4 = \frac{1}{2} \int_{\Omega} v^2 (2\alpha k^2 x + g'_2(x)) dx + (1 - \rho^2) k^2 \int_{\Omega} \frac{\partial u_1}{\partial x} x^2 v \frac{\partial v}{\partial x} dx dy.$$

This term is bounded from below:

$$B_4 \geq \int_{\Omega} (\alpha k^2 - \delta) xv^2 dx dy - \frac{1}{2} |\rho| k \|f'_1\|_{L^\infty(\mathbb{R}_+)} \|v\|_{L^2(\Omega)}^2 \\ - (1 - \rho^2) \left(\epsilon_3 \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 + \frac{k^4}{4\epsilon_3} \left\| x \frac{\partial u_1}{\partial x} \right\|_{L^\infty(\Omega)}^2 \|v\|_{L^2(\Omega)}^2 \right)$$

and we obtain

$$B_4 \geq -\epsilon_2 \|xv\|_{L^2(\Omega)}^2 - (1 - \rho^2) \epsilon_3 \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 \\ - \left(\frac{(\alpha k^2 - \delta)^2}{4\epsilon_2} - \frac{1}{2} |\rho| k \|f'_1\|_{L^\infty(\mathbb{R}_+)} + \frac{k^4}{4\epsilon_3} \left\| x \frac{\partial u_1}{\partial x} \right\|_{L^\infty(\Omega)}^2 \right) \|v\|_{L^2(\Omega)}^2.$$

$$B_5 \geq -\frac{|\rho|}{2} \alpha k \|xv\|_{L^2(\Omega)}^2.$$

$$B_6 \geq \alpha \left(\delta - \frac{\alpha k^2}{2} \right) \|xv\|_{L^2(\Omega)}^2 - \int_{\Omega} \alpha (\delta \sigma_1 + (1 - \rho^2) k^2 \left\| x \frac{\partial u_1}{\partial x} \right\|_{L^\infty(\Omega)}^2) xv^2 dx dy \\ - \alpha |\rho| k \|f_1\|_{L^\infty(\mathbb{R}_+)} \|v\|_{L^2(\Omega)}^2.$$

and we deduce :

$$B_6 \geq \left(\alpha \left(\delta - \frac{\alpha k^2}{2} \right) - \epsilon_4 \right) \|xv\|^2 \\ - \left(\frac{\alpha^2}{4\epsilon_4} (\delta \sigma_1 + (1 - \rho^2) k^2 \left\| x \frac{\partial u_1}{\partial x} \right\|_{L^\infty(\Omega)}^2)^2 + \alpha |\rho| k \|f_1\|_{L^\infty(\mathbb{R}_+)} \right) \|v\|_{L^2(\Omega)}^2.$$

From all these estimates, we get:

$$\hat{b}(v, v) \geq (1 - |\rho|) \frac{k^2}{2} (1 - (1 + |\rho|) \epsilon_3) \left\| x \frac{\partial v}{\partial x} \right\|_{L^2(\Omega)}^2 + \frac{1}{2} (1 - |\rho|) \left\| xy \frac{\partial v}{\partial y} \right\|_{L^2(\Omega)}^2 \\ + \left(\alpha \left(\delta - \frac{\alpha k^2}{2} \right) - \frac{1}{2} - |\rho| \frac{\alpha k}{2} - \epsilon \right) \|xv\|_{L^2(\Omega)}^2 - c \|v\|_{L^2(\Omega)}^2$$

with $\epsilon = \epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4$ and c is a positive constant depending on the different parameters.

$$\text{Let us denote } P(\alpha) = \alpha \left(\delta - \frac{\alpha k^2}{2} \right) - \frac{1}{2} - |\rho| \frac{\alpha k}{2} - \epsilon.$$

This polynomial admits two positive real roots if $k < \frac{2}{3}\delta$ and ϵ small enough. This last condition is generally satisfied in practice ($k = 0.4$, $\delta = 2$). So we can choose $\alpha > 0$ such that $P(\alpha) > 0$. Therefore by choosing ϵ_i , $1 \leq i \leq 4$ small enough, α such that $P(\alpha) > 0$ and if $|\rho| < 1$, we obtain:
 $\hat{b}(v, v) \geq C \|v\|_2^2 - c \|v\|_{L^2(\Omega)}^2$.

We may write problem (3.1) in variational form:

$$(3.1) \quad \begin{cases} \text{Find } \hat{u}_2 \in C(0, T; L^2(\Omega)) \cap L^2(0, T; \hat{V}_2) \\ (\frac{\partial \hat{u}_2}{\partial t}, v) + \hat{b}(\hat{u}_2, v) = 0, \quad \forall v \in V_2 \\ \hat{u}_2(0) = e^{-\alpha x} F \end{cases}.$$

Since $\hat{u}_2(0) \in \hat{V}_2$, by using proposition 3.1, we deduce the following theorem [5]:

Theorem 3.2. *Under the hypotheses of Proposition 3.1, problem (3.1) admits a unique solution*

3.2. Numerical solution.

For the computation of u_2 , we shall use a backward Euler method in time and a finite element method in space [4].

We define a triangulation $\mathcal{T}_h$ of Ω in the following way:

- (x_i) , $0 \leq i \leq N$ is the sequence defined in section (2.1.2).
- Let (y_j) , $0 \leq j \leq M$ ($y_0 = 0$) another increasing sequence.

We denote $k_j = y_j - y_{j-1}$ and we assume that the sequence (k_j) , $1 \leq j \leq M$ is increasing.

The domain $[0, x_N] \times [0, y_M]$ is divided in rectangles $(x_{i-1}, x_i) \times (y_{j-1}, y_j)$, $1 \leq i \leq N$, $1 \leq j \leq M$ and each rectangle is divided in two triangles by the first diagonal. We denote $\mathcal{T}_h^1$ the set of these triangles.

We define $\mathcal{T}_h^2$, $\mathcal{T}_h^3$, K_{NM} by:

$$\mathcal{T}_h^2 = \{K_j/K_j = (x_N, +\infty) \times (y_{j-1}, y_j), j = 1, M\}$$

$$\mathcal{T}_h^3 = \{K_i/K_i = (x_{i-1}, x_i) \times (y_M, +\infty), i = 1, N\}$$

$$K_{NM} = (x_N, +\infty) \times (y_M, +\infty)$$

The triangulation is then defined by: $\mathcal{T}_h = \mathcal{T}_h^1 \cup \mathcal{T}_h^2 \cup \mathcal{T}_h^3 \cup K_{NM}$.

We associate to this triangulation the finite-dimensional space V_{2h} defined by:

$$V_{2h} = \left\{ v_h \in C^0(\Omega) / \forall K \in \mathcal{T}_h^1, v_h|_K \in P_1, \forall K \in \mathcal{T}_h^2, v_h|_K \in P_{y1} \right. \\ \left. \forall K \in \mathcal{T}_h^3, yv_h|_K \in P_{x1}, yv_h|_{K_{NM}} \in P_0 \right\}$$

P_1 is the space of polynomials of degree ≤ 1 in x, y ; P_{y1} is the space of polynomials of degree ≤ 1 in y ; P_{x1} is the space of polynomials of degree ≤ 1 in x ; P_0 is the space of constants.

Let $\alpha > 0$ such that $P(\alpha) > 0$.

If $v_h \in V_{2h}$, it is easily seen that $\hat{v}_h = e^{-\alpha x} v_h \in \hat{V}_2$.

The approximate value of u_2 at the time level t_n will be in V_{2h} .

If $v_h \in V_{2h}$, we denote $\tilde{v}_h = e^{-2\alpha x} v_h$ and we define the bilinear form b on $V_{2h} \times V_{2h}$ by: $\forall u_h, v_h \in V_{2h}$,

$$\begin{aligned} b(u_h, v_h) = & \frac{k^2}{2} \int_{\Omega} \frac{\partial u_h}{\partial x} \frac{\partial}{\partial x} (x^2 \tilde{v}_h) dx dy + \frac{1}{2} \int_{\Omega} x^2 \frac{\partial u_h}{\partial y} \frac{\partial}{\partial y} (y^2 \tilde{v}_h) dx dy \\ & + \rho k \int_{\Omega} x^2 \frac{\partial u_h}{\partial x} \frac{\partial}{\partial y} (y \tilde{v}_h) dx dy - \int_{\Omega} x g_2(x) \frac{\partial u_h}{\partial x} \tilde{v}_h dx dy \\ & - (1 - \rho^2) k^2 \int_{\Omega} x^2 \frac{\partial u_h}{\partial x} \frac{\partial u_h}{\partial x} \tilde{v}_h dx dy. \end{aligned}$$

We have the equality: $b(u_h, v_h) = \hat{b}(\hat{u}_h, \hat{v}_h)$, $\forall u_h, v_h \in V_{2h}$ and we define the approximate value u_{2h}^{n+1} of $u_2(t_{n+1})$ as the solution of the following problem:

$$\begin{aligned} (u_{2h}^{n+1} - u_{2h}^n, \tilde{v}_h)_h + b_h(u_{2h}^{n+1}, v_h) &= 0, \forall v_h \in V_{2h}, \\ u_{2h}^0 &= F_h, \end{aligned}$$

where F_h is the Lagrange interpolate of F in V_{2h} ;

$(u_h, v_h)_h$ is an approximate scalar product in $L^2(\Omega)$ and b_h an approximation of b , obtained by using numerical integration.

In Fig 2, we present the variation of $u_2(T)$ in y (or $b(0)$ in P) for different values of the volatility; the parameter ρ is null; the exercise price E is equal to 1; the others parameters have the same values as in Fig1.




FIGURE 2. $b(0)$ with $T=1$

Fig 3 represents the variation of $u_2(T)$ in y for different values of ρ , the volatility is equal to 0.3.




FIGURE 3. $b(0)$ with $\sigma=0.3$

4. COMPUTATION OF u_3

u_3 is solution of

$$\begin{aligned} & \frac{\partial u_3}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 u_3}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 u_3}{\partial y^2} - \rho k x^2 y \frac{\partial^2 u_3}{\partial x \partial y} - g(x) \frac{\partial u_3}{\partial x} - x f(x) y \frac{\partial u_3}{\partial y} \\ &= (1 - \rho^2) k^2 x^2 \exp(u_1) \left(\frac{\partial u_2}{\partial x} \right)^2, \end{aligned}$$

with the initial condition: $u_3(0) = 0$.

4.1. Existence and uniqueness of the solution. As for the computation of u_2 , we make the change of unknown: $\hat{u}_3 = e^{-\alpha x} u_3$ and $\hat{u}_3$ is solution of:

$$\begin{aligned} & \frac{\partial \hat{u}_3}{\partial t} - \frac{k^2 x^2}{2} \frac{\partial^2 \hat{u}_3}{\partial x^2} - \frac{x^2 y^2}{2} \frac{\partial^2 \hat{u}_3}{\partial y^2} - \rho k x^2 y \frac{\partial^2 \hat{u}_3}{\partial x \partial y} - \frac{\partial \hat{u}_3}{\partial x} (\alpha k^2 x^2 + g(x)) - \frac{\partial \hat{u}_3}{\partial y} (\rho k \alpha x^2 y + x f(x) y) \\ (4.1) \quad & - \hat{u}_3 \left(\frac{\alpha^2 k^2 x^2}{2} + \alpha g(x) \right) = (1 - \rho^2) k^2 x^2 \exp(u_1 - \alpha x) \left(\frac{\partial u_2}{\partial x} \right)^2. \end{aligned}$$

We define the bilinear form c on $\hat{V}_2 \times \hat{V}_2$ by:

$$\begin{aligned}
\forall u, v \in \hat{V}_2, \quad \hat{c}(u, v) = & \frac{k^2}{2} \int_{\Omega} \frac{\partial u}{\partial x} \frac{\partial}{\partial x} (x^2 v) dx dy + \frac{1}{2} \int_{\Omega} x^2 \frac{\partial u}{\partial y} \frac{\partial}{\partial y} (y^2 v) dx dy \\
& + \rho k \int_{\Omega} x^2 \frac{\partial u}{\partial x} \frac{\partial}{\partial y} (y v) dx dy - \int_{\Omega} (\alpha k^2 x^2 + g(x)) \frac{\partial u}{\partial x} v dx dy \\
& - \int_{\Omega} (\rho \alpha k x^2 + x f(x)) y \frac{\partial u}{\partial y} v dx dy - \int_{\Omega} \left(\frac{\alpha^2 k^2 x^2}{2} + \alpha g(x) \right) u v dx dy.
\end{aligned}$$

It is clear that this bilinear form is continue on $\hat{V}_2 \times \hat{V}_2$ and under the same hypotheses as for b , it satisfies: $\forall v \in \hat{V}_2, \quad \hat{c}(v, v) \geq C \|v\|_{\hat{V}_2}^2 - c \|v\|_{L^2(\Omega)}^2$.

We denote $G(x) = (1 - \rho^2) k^2 x^2 e^{u_1} \left(\frac{\partial u_2}{\partial x} \right)^2, \quad x > 0$.

Problem (4.1) may be written in variational form:

$$(4.2) \quad \begin{cases} \text{Find } \hat{u}_3 \in C^2(0, T; L^2(\Omega) \cap L^2(0, T; \hat{V}_2) \\ \left(\frac{\partial \hat{u}_3}{\partial t}, v \right) + \hat{c}(\hat{u}_3, v) = (G e^{-\alpha x}, v) \\ \hat{u}_3(0) = 0 \end{cases}.$$

If we assume that $\frac{\partial u_2}{\partial x} \in C(0, T; L^\infty(\Omega))$, the second member is continue on $\hat{V}_2$ and we get the following theorem:

Theorem 4.1. *If $|\rho| < 1$, $k < \frac{2}{3}\delta$ and α such that $\hat{c}$ satisfies Garding inequality, problem (4.2) admits a unique solution.*

4.2. Numerical solution. To compute u_3 , we use the same method as for the computation of u_2 .

If $v_h \in V_{2h}$, we denote $\tilde{v}_h = e^{-2\alpha x} v_h$ and define the bilinear form on $V_{2h} \times V_{2h}$ by:

$$\begin{aligned}
c(u_h, v_h) = & \frac{k^2}{2} \int_{\Omega} \frac{\partial u_h}{\partial x} \frac{\partial}{\partial x} (x^2 \tilde{v}_h) dx dy + \frac{1}{2} \int_{\Omega} x^2 \frac{\partial u_h}{\partial y} \frac{\partial}{\partial y} (y^2 \tilde{v}_h) dx dy + \rho k \int_{\Omega} x^2 \frac{\partial u_h}{\partial x} \frac{\partial}{\partial y} (y \tilde{v}_h) dx dy \\
& - \int_{\Omega} g(x) \frac{\partial u_h}{\partial x} \tilde{v}_h dx dy - \int_{\Omega} x f(x) y \frac{\partial u_h}{\partial x} \tilde{v}_h dx dy
\end{aligned}$$

and we have the equality: $\forall u_h, v_h \in V_{2h}, \quad c(u_h, v_h) = \hat{c}(\hat{u}_h, \hat{v}_h)$.

The approximate solution u_{3h}^{n+1} at the time level t_{n+1} satisfies:

$$\begin{aligned}
& (u_{3h}^{n+1} - u_{3h}^n, \tilde{v}_h)_h + \Delta t_n c_h(u_{3h}^{n+1}, v_h) = (G, \tilde{v}_h)_h, \quad \forall v_h \in V_{2h}, \\
& u_{3h}^0 = 0.
\end{aligned}$$

where c_h is an approximation of c obtained by using numerical integration.

Fig4 represents the variation of $u_3(T)$ in y (or $c(0)$ in P) for different values of σ . The parameter ρ is equal to 0. (In the case $\rho = 1$ or $\rho = -1$, u_3 is null). The replication error ϵ^* is given by: $\epsilon^* = \sqrt{c(0)}$.




FIGURE 4. $c(0)$ with $T=1$

Knowing u_2 , we can compute the least-cost optimal strategy $\theta^*(0) = \frac{\partial b}{\partial P}(0) + \frac{\rho k}{P} \frac{\partial b}{\partial \sigma}(0) = \frac{\partial u_2}{\partial y}(T) + \frac{\rho k}{y} \frac{\partial u_2}{\partial y}(T)$.

Fig 5 represents the variation of θ^* in P for different values of σ ; the parameter ρ is equal to 0.

REFERENCES

- [1] D. Bertsimas, L. Kogan, and A.W. Lo. Hedging derivative securities and incomplete markets: An ϵ -arbitrage approach. *Operations Research*, 49(3):372–397, 2001.
- [2] F. Black and M. Scholes. Pricing of options and corporate liabilities. *J. of Political Econom.*, 81:637–654, 1973.
- [3] P. Bolley and J. Camus J. Quelques résultats sur les espaces de Sobolev avec poids. *Publications de Séminaires Mathématiques de Rennes*, 1:1–69, 1969.
- [4] P.G. Ciarlet. *The Finite Element Method for Elliptic Problems*. North-Holland Publishing Company, 1978.
- [5] J.L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications*, volume 1. Dunod, Paris, 1968.
- [6] R.D. Richtmyer and K.W. Morton. *Difference Methods for Initial-Value Problems*. 2nd ed, Wiley-Interscience, New York, 1967.
- [7] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura. Appl.*, 146:65–96, 1987.




FIGURE 5. Optimal strategy

UNIVERSITE BORDEAUX1, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR 5251,351,COURS DE LA
LIBÉRATION, 33405, TALENCE CEDEX
E-mail address: Marie-Noelle.Leroux@math.u-bordeaux1.fr